arXiv:1509.05542v1 [math.GN] 18 Sep 2015

ON THE SEQUENTIAL CLOSURE OF THE SET OF CONTINUOUS FUNCTIONS IN THE SPACE OF SEPARATELY CONTINUOUS FUNCTIONS

TARAS BANAKH

Abstract. For separable metrizable spaces X, Y and a metrizable topological group Z by $S(X\times Y,Z)$ we denote the space of all separately continuous functions $f:X\times Y\to Z$ endowed with the topology of layer-wise uniform convergence, generated by the subbase consisting of the sets $[K_X\times K_Y,U]=\{f\in S(X\times Y,Z):f(K_X\times K_Y)\subset U\}$, where U is an open subset of Z and $K_X\subset X$, $K_Y\subset Y$ are compact sets one of which is a singleton. We prove that every separately continuous function $f:X\times Y\to Z$ with zero-dimensional image $f(X\times Y)$ is a limit of a sequence of jointly continuous functions in the topology of layer-wise uniform convergence.

In this paper we study the sequential closure $\bar C^s(X\times Y,Z)$ of the set $C(X\times Y,Z)$ of continuous functions in the space $S(X\times Y,Z)$ of separately continuous functions $f:X\times Y\to Z$ defined on the product of topological spaces X,Y with values in a topological space Z. The space $S(X\times Y,Z)$ is endowed with the topology of layer-wise compact-open topology, generated by the subbase consisting of the sets

$$[K_X\times K_Y,U]=\{f\in S(X\times Y,Z):f(K_X\times K_Y)\subset U\},$$

where $U\subset Z$ is an open set and $K_X\subset X$, $K_Y\subset Y$ are compact subsets one of which is a singleton. Observe that a function sequence $(f_n)_{n\in\omega}$ converges to a function $f_\infty:X\times Y\to Z$ in the space $S(X\times Y,Z)$ if and only if for any points $x\in X$, $y\in Y$ the function sequence $\big(f_n(x,\cdot)\big)_{n\in\omega}$ converges to $f_\infty(x,\cdot)$ in the space $C(Y,Z)$ and the function sequence $\big(f_n(\cdot,y)\big)_{n\in\omega}$ converges to $f_\infty(\cdot,y)$ in the function space $C(X,Z)$ (endowed with the compact-open topology).

In case of compact spaces X,Y and the real line $Z=\mathbb R$ the space $S(X\times Y,Z)$ was studied by H.A. Voloshyn, V.K. Maslyuchenko and O.V. Maslyuchenko in the papers [1]–[5]. One of the problems addressed in these papers was the problem of description of the sequential closure $\bar C^s(X\times Y,Z)$ of the set $C(X\times Y,Z)$ of jointly continuous functions in the space $S(X\times Y,Z)$. By the *sequential closure* $\bar A^s$ of a set $A\subset X$ in a topological space T we understand the set of all limit points $\lim_{n\to\infty}a_n$ of sequences $\{a_n\}_{n\in\omega}\subset A$, convergent in the space T. The Tietze-Urysohn Extension Theorem [8, 2.1.8] implies that for any Tychonoff spaces X,Y the set $C(X\times Y,\mathbb R)$ is dense in $S(X\times Y,\mathbb R)$, which motivates the following question.

Problem 1. *Is $\bar C^s(X\times Y,\mathbb R)=S(X\times Y,\mathbb R)$ for any (zero-dimensional) compact metrizable spaces X,Y?*

In [3] Voloshyn and Maslyuchenko proved that for metrizable compact spaces X,Y the set $\bar C^s(X\times Y,\mathbb R)$ contains all separately continuous functions $f:X\times Y\to\mathbb R$ whose discontinuity point set $D(f)\subset X\times Y$ has countable projection on X or Y.

In this paper we shall prove that for separable metrizable spaces X,Y and a metrizable topological group Z the set $\bar C^s(X\times Y,Z)$ contains all separately continuous functions $f:X\times Y\to Z$ with zero-dimensional image $f(X\times Y)\subset Z$.

We start with the following useful fact.

Theorem 1. *For separable metrizable spaces X,Y and a topological space Z the set $\bar C^s(X\times Y,Z)$ contains all separately continuous functions $f:X\times Y\to Z$ with discrete image $f(X\times Y)\subset Z$.*

Proof. Fix a separately continuous function $f:X\times Y\to Z$ with discrete image $f(X\times Y)\subset Z$. We lose no generality assuming that $Z=f(X\times Y)$. The separate continuity of f implies that for any countable dense subsets $D_X\subset X$ and $D_Y\subset Y$ the set $f(D_X\times D_Y)$ is dense in the discrete space $Z=f(X\times Y)$, which implies that $Z=f(D_X\times D_Y)$ is countable and hence can be written as the countable union $Z=\bigcup_{n\in\omega}Z_n$ of a non-decreasing sequence $(Z_n)_{n\in\omega}$ of finite sets.

The second-countable spaces X,Y have countable bases $\{U_n\}_{n\in\omega}$ and $\{V_n\}_{n\in\omega}$ of the topology.

1991 *Mathematics Subject Classification.* 54C08, 54C05, 26B05.

For any point $z\in Z$ and a number $k\in\omega$ consider the sets

$$X(z,k)=\{x\in X:\{x\}\times V_k\subset f^{-1}(z)\}=\{x\in X:\{x\}\times\overline{V}_k\subset f^{-1}(z)\},$$
$$Y(z,k)=\{y\in Y:U_k\times\{y\}\subset f^{-1}(z)\}=\{y\in Y:\overline{U}_k\times\{y\}\subset f^{-1}(z)\}.$$

We claim that the set $X(z,k)$ is closed in X. Indeed, if $x\in X\setminus X(z,k)$, then $f(x,v)\ne z$ for some $v\in V_k$. The separate continuity of f yields a neighborhood $O_x\subset X$ of x such that $f(O_x\times\{v\})\subset Z\setminus\{z\}$, which implies that $O_x\cap X(z,k)=\emptyset$. By analogy we can prove that the sets $Y(z,k)$ are closed in Y.

It follows that for any number $n\in\omega$ the set

$$XY(z,n)=\bigcup_{k\le n}(X(z,k)\times\overline{V}_k)\cup(\overline{U}_k\times Y(z,k))$$

is closed in $X\times Y$ and is contained in the preimage $f^{-1}(z)$. Consequently, the family of closed sets $\big(XY(z,n)\big)_{z\in Z}$ is disjoint. Using the normality of the product $X\times Y$, we can construct a continuous function $g_n:X\times Y\to Z$ such that $XY(z,n)\subset g_n^{-1}(z)$ for all $z\in Z_n$.

It remains to check that the sequence $(g_n)_{n\in\omega}$ converges to f in the space $S(X\times Y,Z)$. Given a neighborhood $O_f\subset S(X\times Y,Z)$, it suffices to find a number $m\in\omega$ such that $g_n\in O_f$ for all $n\ge m$. We lose no generality assuming that O_f is a subbasic neighborhood of the form $[K_X\times K_Y,W]=\{g\in S(X\times Y,Z):g(K_X\times K_Y)\subset W\}$ for some some set $W\subset Z$ and some compacts sets $K_X\subset X$, $K_Y\subset Y$, one of which is a singleton. Since the set $f(K_X\times K_Y)\subset W\subset Z$ is compact and hence finite in the discrete space Z, we can replace W by the image $f(K_X\times K_Y)$ and assume that $W=f(K_X\times K_Y)$ is a finite (open) set in X.

First we consider the case of singleton set $K_X=\{x\}$. Consider the continuous map $f_x:Y\to Z$, $f_x:y\mapsto f(x,y)$, and observe that the set $f_x^{-1}(W)$ is closed-and-open in Y and contains the compact set K_Y. For every $z\in W$ find a finite subset $F_z\subset\omega$ such that $K_Y\cap f_x^{-1}(z)\subset\bigcup_{k\in F_z}V_k\subset f_x^{-1}(z)$. Choose a number $m\in\omega$ such that $W\subset Z_m$ and $\bigcup_{z\in W}F_z\subset\{0,\dots,m\}$. Then for every $n\ge m$ and $z\in W$ we get $\{x\}\times(K_Y\cap f_x^{-1}(z))\subset XY(z,n)\subset g_n^{-1}(z)$ and hence

$$K_X\times K_Y=\{x\}\times K_Y\subset\bigcup_{z\in W}\{x\}\times f_x^{-1}(z)\subset\bigcup_{z\in W}g_n^{-1}(z)=g_n^{-1}(W),$$

which implies $g_n\in O_f=[K_X\times K_Y,W]$ for all $n\ge m$.

By analogy we can treat the case of singleton set K_Y. □

Recall that a topological space Z is *zero-dimensional* if closed-and-open sets form a base of the topology of the space Z.

Theorem 2. *For separable metrizable spaces X,Y and a metrizable topological group G, the set $\bar{C}^s(X\times Y,Z)$ contains all separately continuous functions $f:X\times Y\to G$ with zero-dimensional image $f(X\times Y)\subset G$.*

Proof. Fix any separately continuous function $f:X\times Y\to G$ with a zero-dimensional image $f(X\times Y)$. By Birkhof-Kakutani Theorem [6, 3.3.12], the topology of the metrizable topological group G is generated by a bounded left-invariant metric d. Multiplying d by a small positive constant, we can assume that $\mathrm{diam}(G)=\sup\{d(x,y):x,y\in G\}\le\frac12$. For a positive real ε denote by $B[\varepsilon]=\{g\in G:d(g,1_G)\le\varepsilon\}$ the closed ε-ball centered at the unit 1_G of the group G. For every $n\in\omega$ choose a maximal subset $E_n\subset B[\frac{1}{2^{n-1}}]$, which is $\frac{1}{2^{n+1}}$-*separated* in the sense that $d(x,y)\ge\frac{1}{2^{n+1}}$ for any distinct points $x,y\in E_n$. It is clear that the set E_n is closed and discrete in G.

The separability of the spaces X,Y and the separate continuity of $f:X\times Y\to G$ implies the separability of the image $f(X\times Y)$. Since the metrizable space $Z=f(X\times Y)$ is separable and zero-dimensional, it has covering dimension $\dim(Z)=0$, which allows us to choose a sequence $(\mathcal{W}_n)_{n\in\omega}$ of disjoint open covers of the space Z by sets of diameter $\le\frac{1}{2^{n+1}}$. Replacing each cover $\mathcal{W}_n$ by the finer cover $\{W\cap W':W\in\mathcal{W}_n,\ W'\in\mathcal{W}_{n+1}\}$, we can assume that for every $n\in\omega$ the cover $\mathcal{W}_{n+1}$ refines the cover $\mathcal{W}_n$. Since $\mathrm{diam}(Z)\le\frac12$, we can put $\mathcal{W}_0=\{Z\}$.

Inductively we shall construct a sequence $(r_n)_{n\in\omega}$ of continuous maps $r_n:Z\to G$ such that $r_0(Z)=\{1_G\}$ and for every $n\in\omega$ the following conditions are satisfied:

- (1_n) for each set $W\in\mathcal{W}_n$ the image $r_n(W)$ is a singleton;
- (2_n) $\sup_{z\in Z}d(r_n(z),z)\le 2^{-n}$;
- (3_n) $r_n(z)\in r_{n-1}(z)\cdot E_{n+1}$ for every $z\in Z$.

We start the inductive construction putting $r_0:Z\to\{1_G\}\subset G$. Assume that for some $n\in\omega$ we have constructed a map $r_n:Z\to G$ satisfying the conditions (1_n) and (2_n). For every $W\in\mathcal{W}_n$ denote by g_W the unique point of the set $r_n(W)$. Consider the subfamily $\mathcal{W}_{n+1}(W)=\{W'\in\mathcal{W}_{n+1}:W'\subset W\}$ and for every set $W' \in \mathcal{W}_{n+1}(W)$ choose a point $z_{W'} \in W'$. Observe that $d(z_{W'}, g_W) = d(z_{W'}, r_n(z_{W'})) \le \frac{1}{2^n}$ and hence $g_W^{-1} z_{W'} \in B[\frac{1}{2^n}]$. By the maximality of the $\frac{1}{2^{n+2}}$-separated set $E_{n+1} \subset B[\frac{1}{2^n}]$, there exists a point $\varepsilon_{W'} \in E_{n+2}$ such that $d(\varepsilon_{W'}, g_W^{-1} z_{W'}) < \frac{1}{2^{n+2}}$. Consider the point $g_{W'} = g_W \cdot \varepsilon_{W'}$ and observe that

$$d(g_{W'}, z_{W'}) = d(g_W \cdot \varepsilon_{W'}, z_{W'}) = d(\varepsilon_{W'}, g_W^{-1} z_{W'}) \le \frac{1}{2^{n+2}}.$$

Define a function $r_{n+1} : Z \to G$ assigning to each point $z \in Z$ the point $g_{W'}$ where $W' \in \mathcal{W}_{n+1}$ is a unique set containing the point z. It is clear that the function r_{n+1} is continuous and satisfies the condition (1_{n+1}). To check the condition (2_{n+1}), take any point $z \in Z$ and find a unique set $W' \in \mathcal{W}_{n+1}$ containing the point z and a unique set $W \in \mathcal{W}_n$ containing the set W'. Observe that

$$d(r_{n+1}(z), z) = d(g_{W'}, z) \le d(g_{W'}, z_{W'}) + d(z_{W'}, z) \le 2^{-n-2} + \operatorname{diam}(W') \le 2^{-n-2} + 2^{-n-2} = 2^{-n-1}.$$

To verify the condition (3_{n+1}), take any point $z \in Z$ and find unique sets $W \in \mathcal{W}_n$ and $W' \in \mathcal{W}_{n+1}$ containing the point z. Taking into account that the cover $\mathcal{W}_{n+1}$ refines the disjoint cover $\mathcal{W}_n$, we conclude that $W' \subset W$, which implies $r_{n+1}(z) = g_{W'} = g_W \cdot \varepsilon_{W'} = r_n(z) \cdot \varepsilon_{W'} \subset r_n(z) \cdot E_{n+1}$. This completes the inductive construction of the function sequence $(r_n)_{n \in \omega}$.

The continuity of r_n and the separate continuity of f implies that the composition $f_n = r_n \circ f : X \times Y \to G$ is separately continuous and

$$\sup_{(x,y) \in X \times Y} d(f_n(x,y), f(x,y)) \le \sup_{z \in Z} d(r_n(z), z) \le 2^{-n},$$

which means that the function sequence $(f_n)_{n \in \omega}$ converges uniformly to f.

The continuity of the group operations on Z and the separate continuity of the functions f_n, f_{n+1} imply that the function $g_n = f_n^{-1} f_{n+1} : X \times Y \to G$, $g_n : (x,y) \mapsto f_n^{-1}(x,y) \cdot f_{n+1}(x,y)$, is separately continuous. By the condition (3_{n+1}), we get $g_n(x,y) = f_n^{-1}(x,y) \cdot f_{n+1}(x,y) = r_n(f(x,y))^{-1} \cdot r_{n+1}(f(x,y)) \in E_{n+1}$, which means that the image $g_n(X \times Y)$ is discrete. By Theorem 1, for the separately continuous function $g_n : X \times Y \to E_{n+1}$ there is a sequence of continuous functions $(g_{n,m} : X \times Y \to E_{n+1})_{m \in \omega}$, convergent to g_n in the function space $S(X \times Y, Z)$.

For any numbers $n, m \in \omega$ consider the continuous function

$$f_{n,m} = g_{0,m} \cdot g_{1,m} \cdots g_{n,m} : X \times Y \to G.$$

Taking into account that $\lim_{m \to \infty} g_{n,m} = g_n$ for all n, we conclude that $\lim_{m \to \infty} f_{n,m} = g_0 \cdots g_n = f_n$ for all $n \in \omega$. It remains to check that $\lim_{n \to \infty} f_{n,n} = f$ in the space $S(X \times Y, G)$.

Fix any neighborhood O_f of the function f in the space $S(X \times Y)$. We lose no generality assuming that O_f has subbasic form $[K_X \times K_Y, W] = \{g \in S(X \times Y, G) : f(K_X \times K_Y) \subset W\}$ for some open set $W \subset G$ and some compact sets $K_X \subset X$, $K_Y \subset Y$ one of which is a singleton. Using the compactness of the set $f(K_X \times K_Y) \subset W$, find a number $l \in \omega$ such that $f(K_X \times K_Y) \cdot B[\frac{4}{2^l}] \subset W$. By the convergence of the sequence $(f_{l,n})_{n \in \omega}$ to the function f_l in the space $S(X \times Y, G)$, there is a number $m \ge l$ such that $\sup_{(x,y) \in K_X \times K_Y} d(f_{l,n}(x,y), f_l(x,y)) \le 2^{-l}$ for all $n \ge m$. We claim that $\sup_{(x,y) \in K_X \times K_Y} d(f(x,y), f_{n,n}(x,y)) \le 2^{-l}$ for all $n \ge m$. Take and number $n \ge m$ and a pair $z = (x,y) \in K_X \times K_Y$. Observe that $g_{l+1,n}(z) \cdots g_{n,n}(z) \in E_{l+2} \cdots E_{n+1} \subset B[\frac{1}{2^{l+1}}] \cdots B[\frac{1}{2^n}] \subset B[\frac{1}{2^l}]$. Then

$$\begin{aligned} d(f(z), f_{n,n}(z)) &\le d(f(z), f_l(z)) + d(f_l(z), f_{l,n}(z)) + d(f_{l,n}(z), f_{n,n}(z)) \le \\ &\le 2^{-l} + 2^{-l} + d(g_{0,n}(z) \cdots g_{l,n}(z), g_{0,n}(z) \cdots g_{n,n}(z)) = \\ &= 2^{-l+1} + d(1_G, g_{l+1,n}(z) \cdots g_{n,n}(z)) \le 2^{-l+1} + 2^{-l} < 2^{-l+2} \end{aligned}$$

and hence $f_{n,n}(z) \in W$ according to the choice of l. □

Now we find conditions on a topological group Z guaranteeing that the sequential closure $\bar{C}^s(X \times Y, Z)$ of the set $C(X \times Y, Z)$ of continuous functions in the space $S(X \times Y, Z)$ is closed in a topology of uniform convergence on $S(X \times Y, Z)$.

In fact, the space $S(X \times Y, Z)$ carries four natural topologies of uniform convergence induced by four canonical uniformities on the topological group Z. For describing these topologies, for any function $f \in S(X \times Y, Z)$ and any neighborhood $U \subset Z$ of the unit in the group Z consider the sets

$$B_l[f,U] = \{g \in S(X \times Y, Z) : \forall (x,y) \in X \times Y \;\; g(x,y) \in f(x,y) \cdot U\},$$
$$B_r[f,U] = \{g \in S(X \times Y, Z) : \forall (x,y) \in X \times Y \;\; g(x,y) \in U \cdot f(x,y)\},$$
$$B_{lr}[f,U] = B_l[f,U] \cap B_r[f,U] = \{g \in S(X \times Y, Z) : \forall (x,y) \in X \times Y \;\; g(x,y) \in U \cdot f(x,y) \cap f(x,y) \cdot U\},$$
$$B_{rl}[f,U] = \{g \in S(X \times Y, Z) : \forall (x,y) \in X \times Y \;\; g(x,y) \in U \cdot f(x,y) \cdot U\}.$$

We define a subset $W \subset S(X \times Y, Z)$ to be τ_s-*open* for $s \in \{l, r, lr, rl\}$ if for any function $f \in W$ there is a neighborhood $U \subset Z$ of the unit 1_Z in Z such that $B_s[f,U] \subset W$. The family τ_s of all τ_s-open subsets of $S(X \times Y, Z)$ is a topology on $S(X \times Y, Z)$. It is clear that $\tau_{rl} \subset \tau_l \cap \tau_r$ and $\tau_l \cup \tau_r \subset \tau_{lr}$. We define a subset $F \subset S(X \times Y, Z)$ to be $\tau_l \cap \tau_r$-*closed* if its complement in $S(X \times Y, Z)$ belongs to the topology $\tau_l \cap \tau_r$.

For topological spaces X, Y, Z by $S_0(X \times Y, Z)$ we denote the set of all separately continuous functions $f : X \times Y \to Z$ with zero-dimensional image $f(X \times Y)$. If Z is a topological group, then by $\bar{S}^u_0(X \times Y, Z)$ we shall denote the closure of the set $S_0(X \times Y, Z)$ in the topology $\tau_l \cap \tau_r$ on the space $S(X \times Y, Z)$. By Theorem 2, $S_0(X \times Y, Z) \subset \bar{C}^s(X \times Y, Z)$ for any metrizable compact spaces X, Y and any metrizable topological group Z. In the following theorem we give conditions on the group Z guaranteeing that $\bar{S}^u_0(X \times Y, Z) \subset \bar{C}^s(X \times Y, Z)$.

We shall say that a topological space Z is an *absolute extensor* for a topological space X if any continuous map $f : A \to Z$ defined on a closed subset $A \subset X$ can be extended to a continuous map $\bar{f} : X \to Z$.

Theorem 3. *Let X, Y be separable metrizable spaces and Z be a metrizable topological group possessing a neighborhood base at the unit 1_Z consisting of absolute extensors for the space $X \times Y$. Then the set $\bar{C}^s(X \times Y, Z)$ is closed in the topology $\tau_l \cap \tau_r$ on $S(X \times Y, Z)$. Consequently, $\bar{S}^u_0(X \times Y, Z) \subset \bar{C}^s(X \times Y, Z)$.*

Proof. It suffices to check that the set $\bar{C}^s(X \times Y, Z)$ is closed in the topologies τ_l and τ_r on the space $S(X \times Y, Z)$. First we prove that the set $\bar{C}^s(X \times Y, Z)$ is τ_l-closed. Fix a left-invariant metric $d \le 1$ generating the topology of the metrizable topological group Z. Put $W_{-1} = W_0 = Z$ and for every $n > 0$ choose a neighborhood $W_n \subset \{z \in Z : d(1_Z, z) \le 2^{-n}\}$ of the unit 1_Z, which is an absolute extensor for the space $X \times Y$. Replacing the sequence $(W_n)_{n \in \omega}$ by a suitable subsequence, we can assume that $W_n^{-1} W_n W_n \subset W_{n-1}$ for all $n > 0$.

Take any function $f \in S(X \times Y, Z)$ that belongs to the τ_l-closure of the set $\bar{C}^s(X \times Y, Z)$ in $S(X \times Y, Z)$. For every $n \in \omega$ choose a function $f_n \in \bar{C}^s(X \times Y, Z)$ such that $f_n(x,y) \in f(x,y) \cdot W_{n+1}$ for all $(x,y) \in X \times Y$. Since $W_0 = Z$ we can put $f_0 : X \times Y \to \{1_Z\}$ be the constant function. For every $n \in \omega$ consider the function $g_n = f_n^{-1} \cdot f_{n+1} : X \times Y \to Z$ and observe that for every $(x,y) \in X \times Y$ we get $g_n(x,y) = f_n^{-1}(x,y) \cdot f_{n+1}(x,y) \subset W_n^{-1} \cdot f(x,y)^{-1} \cdot f(x,y) \cdot W_{n+1} = W_n^{-1} W_{n+1} \subset \overline{W_n^{-1} W_{n+1} W_{n+2}} \subset W_n^{-1} W_{n+1} W_{n+2} W_{n+3} \subset W_{n-1}$.

The continuity of the group operations on Z implies that the set $\bar{C}^s(X \times Y, Z)$ is a group. So, $g_n \in \bar{C}^s(X \times Y, Z)$ and we can choose a sequence $(g_{n,m})_{m \in \omega}$ of continuous functions $g_{n,m} : X \times Y \to G$, which converges to the function g_n in the space $S(X \times Y, Z)$. The functions $g_{n,m}$ can be modified so that $g_{n,m}(X \times Y) \subset W_{n-1}$. Since $W_{-1} = W_0 = Z$, this is trivial for $n \le 1$. Put $\bar{g}_{k,m} = g_{k,m}$ for $k \le 1$, $m \in \omega$. Fix any number $n > 1$. By the continuity of the function $g_{n,m}$, the set $F_{n,m} = g_{n,m}^{-1}(\overline{W_n^{-1} W_{n+1} W_{n+2}})$ is closed in $X \times Y$. Since W_{n-1} is an absolute extensor for the space $X \times Y$, the function $g_{n,m} | F_{n,m}$ can be extended to a continuous function $\bar{g}_{n,m} : X \times Y \to W_{n-1}$. We claim that the function sequence $(\bar{g}_{n,m})_{m \in \omega}$ converges to g_n. It suffices to check that for any sub-basic neighborhood $O(g_n) = [K_X \times K_Y, W]$ of the function g_n there exists a number $k \in \omega$ such that $\bar{g}_{n,m} \in O(g_n)$ for all $m \ge k$. Here $W \subset Z$ is an open set and $K_X \subset X$, $K_Y \subset Y$ are compact subsets one of which is a singleton. Using the compactness of the set $g_n(K_X \times K_Y) \subset W$, find a number $l \ge n+2$ such that $g_n(K_X \times K_Y) \cdot W_l \subset W$. The convergence of the function sequence $(g_{n,m})_{m \in \omega}$ to g_n yields a number $k \in \omega$ such that $g_{n,m}(K_X \times K_Y) \subset g_n(K_X \times K_Y) \cdot W_l \subset W_n^{-1} W_{n+1} W_l \subset \overline{W_n^{-1} W_{n+1} W_{n+2}}$ for all $m \ge k$. For such numbers m we get the inclusion $K_X \times K_Y \subset F_{n,m}$ implying $\bar{g}_{n,m}(K_X \times K_Y) = g_{n,m}(K_X \times K_Y) \subset g_n(K_X \times K_Y) \cdot W_l \subset W$. Then $\bar{g}_{n,m} \in [K_X \times K_Y, W]$, which means that the sequence $(\bar{g}_{n,m})_{m \in \omega}$ converges to g_n in the space $S(X \times Y, Z)$.

For any $n, m \in \omega$ consider the continuous function $f_{n,m} = g_{0,m} \cdots g_{n,m} : X \times Y \to Z$ and observe that $\lim_{m \to \infty} f_{n,m} = g_0 \cdots g_n = f_{n+1}$ for all $n \in \omega$. Repeating the argument of the proof of Theorem 2, we can show that the function sequence $(f_{n,n})_{n \in \omega}$ converges to f in the space $S(X \times Y, Z)$. Then $f \in \bar{C}^s(X \times Y, Z)$ and hence the set $\bar{C}^s(X \times Y, Z)$ is τ_l-closed in $S(X \times Y, Z)$. By analogy we can prove that this set is τ_r-closed.

The $(\tau_l \cap \tau_r)$-closedness of the set $\bar{C}^s(X \times Y, Z)$ and the inclusion $S_0(X \times Y, Z) \subset \bar{C}^s(X \times Y, Z)$ (proved in Theorem 2) implies the inclusion $\bar{S}^u_0(X \times Y, Z) \subset \bar{C}^s(X \times Y, Z)$. ☐

Remark 1. For the interval $X = Y = [0,1]$ the closedness of the set $\bar{C}^s(X \times Y, \mathbb{R})$ in the topology of uniform convergence on $S(X \times Y, \mathbb{R})$ was proved in [4].

Theorem 3 has two corollaries.

Corollary 1. *For zero-dimensional separable metrizable spaces X, Y and any metrizable topological group Z the set $\bar{C}^s(X \times Y, Z)$ is $(\tau_l \cap \tau_r)$-closed in $S(X \times Y, Z)$ and hence $\bar{S}^u_0(X \times Y, Z) \subset \bar{C}^s(X \times Y, Z)$.*

Proof. The zero-dimensionality of the space $X \times Y$ implies that each closed subset of $X \times Y$ is a retract of the space $X \times Y$. So, each topological space is an absolute extensor for the space $X \times Y$, which allows us to apply Theorem 3 and complete the proof. ☐

Corollary 2. *For separable metrizable spaces X, Y and a locally convex linear metric space Z the set $\bar{C}^s(X \times Y, Z)$ is $(\tau_l \cap \tau_r)$-closed in $S(X \times Y, Z)$ and hence $\bar{S}^u_0(X \times Y, Z) \subset \bar{C}^s(X \times Y, Z)$.*

Proof. By the Dugundji Theorem [7], any convex subset of a locally convex space is an absolute extensor for any metrizable space. Consequently, the locally convex space Z has a neighborhood base of (convex) neighborhoods of zero which are absolute extensors for the space $X \times Y$. This allows us to apply Theorem 3 and complete the proof of the corollary. ☐

Observe that an affirmative answer to the "zero-dimensional" part of Problem 1 can be derived from Corollary 1 and the affirmative answer to the following problem.

Problem 2. *Let X, Y be zero-dimensional compact spaces. Is the set $S_0(X \times Y, \mathbb{R})$ of separately continuous functions $f : X \times Y \to \mathbb{R}$ with zero-dimensional image $f(X \times Y)$ dense in the space $S(X \times Y, \mathbb{R})$ endowed with the topology of uniform convergence $\tau_l = \tau_r$?*

This problem can be equivalently reformulated in more elementary terms.

Problem 3. *Let $f : X \times Y \to \mathbb{R}$ be a separately continuous function defined on the product of two Cantor sets $X = Y = \{0, 1\}^\omega$. Is there a separately continuous function $g : X \times Y \to \mathbb{R}$ with (countable) zero-dimensional image $g(X \times Y)$ such that $\sup_{(x,y)\in X\times Y} |f(x, y) - g(x, y)| \le 1$?*

REFERENCES

[1] H.A. Voloshyn, V.K. Maslyuchenko, *A topologization of the space of separately continuous functions*, Carpathian Math. J. **5**:2 (2013), 199–207 (in Ukrainian).

[2] H.A. Voloshyn, V.K. Maslyuchenko, *On a linear interpolation of vector-valued functions and its applications*, Mat. Stud. **42**:2 (2014), 129–133 (in Ukrainian).

[3] H.A. Voloshyn, V.K. Maslyuchenko, *The sequential closure of the space of jointly continuous functions in the space of separately continuous functions*, Ukr. Mat. Zh. (submitted; in Ukrainian).

[4] H.A. Voloshyn, V.K. Maslyuchenko, O.V. Maslyuchenko, *On layer-wise uniform approximation of separately continuous functions by polynomials*, Mat. Bull. Shevchenko Sci. Soc. **10** (2013), 135–158 (in Ukrainian).

[5] H.A. Voloshyn, V.K. Maslyuchenko, O.V. Maslyuchenko, *Embedding the space of separately continuous functions into a product of Banach spaces and its barreledness*, Mat. Bull. Shevchenko Sci. Soc. **11** (2014), 36–50 (in Ukrainian).

[6] A. Arhangel'skii, M. Tkachenko, *Topological groups and related structures*, Atlantis Studies in Mathematics, 1. Atlantis Press, Paris; World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2008.

[7] J. Dugundji, *An extension of Tietze's theorem*, Pacific J. Math. **1** (1951), 353–367.

[8] R. Engelking, *General Topology*, Heldermann Verlag, Berlin, 1989.

E-mail address: `t.o.banakh@gmail.com`

IVAN FRANKO NATIONAL UNIVERSITY OF LVIV (UKRAINE) AND JAN KOCHANOWSKI UNIVERSITY IN KIELCE (POLAND)